PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5:		(11) International Publication Number: WO 93/23544
C12N 15/49, 9/10, 15/62 A61K 39/21	A1	(43) International Publication Date: 25 November 1993 (25.11.93)
(21) International Application Number: PCT/GB (22) International Filing Date: 14 May 1993		& Fairweather, 19 Royal Exchange So. Glasgow Gi
(30) Priority data: 9210337.3 14 May 1992 (14.05.92)		(81) Designated States: AU, CA, JP, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).
 (71) Applicant (for all designated States except US): TI VERSITY COURT OF THE UNIVERSITY OF GOW [GB/GB]; 2 The Square, University Aven gow G12 8QQ (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): SPIBEY, Norm GB]; 51 Speirs Road, Bearsden, Glasgow G61 2: JARRETT, James, Oswald [GB/GB]; 4 Drumonue, Milngavie, Glasgow G62 8NA (GB). 	F GLA	S-Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.
(54) Title: FUSION PROTEIN FROM FELINE LEU	JKAEN	IIA VIRUS GP70 AS VACCINE
	of ti	ne variable regions in the FeLV gp70 gene
1		
gp 70		p15 eny
57) Abstract		

(5

A vaccine against feline leukaemia virus (FeLV) comprises a protein formed of two proteins derived from FeLV gp70 which are not adjacent in the native protein, particularly VR1 and VR5 or effective fragments thereof. VR5 is preferably attached to the carboxy terminus of VR1. A co-protein such as GST or β-galactosidase may be attached to the amino terminus of the fused protein, such as to stabilise and solubilise the fused protein. Attachments to the carboxy terminus may reduce the immunogenicism.

48 30

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
88	Barbados	CB	United Kingdom	NL.	Netherlands
BE	Belgium	GN	Guinca	NO	Norway
8F	Burkina Faso	GR	Grane	NZ	New Zealand
BC	Bulgaria	HU	Hungary	PL.	Poland
B.J	Benin	1E	Ireland	PT	Portugal
BR	Brazil	ıπ	lialy	RO	Portugai Romania
CA	Canada	JP	Japan -	RU	Russian Federation
œ	Central African Republic	KP	Democratic People's Republic	SD	Swian Pederation
œ	Congo		of Konsa	SE	·
CH	Switzerland	KR	Republic of Korea	SK	Sweden Street Breeze Miles
a	Côte d'ivoire	KZ	Kwalistan	SN	Slovak Republic
CM	Cameroon	ü	Liechtenstein	SU	Sunegal
cs	Cochesionalia .	LK	Sri Lanka		Soviet Union
CZ	Czech Republic	LU	Lanembourg	TD	Chad
DE	Germany	MC	Monago	TG	Togo
DK	Dunmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML.		us	United States of America
FI	Finland	MIL.	Mali	VN	Viet Nam
	* *************************************	MM	Mongolia		

FUSION PROTEIN FROM FELINE LEUKAEMIA VIRUS GP70 AS VACCINE

FIELD OF THE INVENTION

The present invention relates to an immunogenic fused recombinant protein derived from feline leukaemia virus (FeLV), and to a protective vaccine comprising the immunogenic protein.

BACKGROUND OF THE INVENTION

FeLV is a retrovirus of the sub-family oncovirinae.

The virus has a simple genomic organisation with three genes, gag, pol and env, encoding viral proteins.

FeLV occurs in three subgroups, A, B or C, which differ in their env genes. FELV-A is present in every isolate and is efficiently transmitted between cats in nature. Viruses of the other two subgroups arise as variants of FeLV-A in individual cats by recombination of endogenous FeLV env genes (FeLV-B) or mutation within the env gene (FeLV-C). Hence the occurrence of subgroups B or C is dependent upon natural infection of cats with FeLV-A.

FeLV is responsible for several fatal diseases in domestic cats including lymphoid and myeloid leukaemias, fibrosarcomas, anaemia, immunodeficiency, enteritis and reproductive failure. The virus is common in nature: a recent survey in the U.K. indicated that the overall

prevalance of infecti n is appr ximately 15% of sick cats and 6% in healthy cats. These cats have a persistent infection with viraemia and are a source of infection for susceptible animals. FeLV is transmitted either by contact through the transfer of infectious saliva or prenatally across the placenta.

Following natural or experimental exposure of cats to FeLV, there are two possible outcomes. Cats either develop a persistent, life-long infection or recover. Persistently infected cats have a viraemia while recovered cats have no virus in their blood. Some cats which recover may harbour a latent infection in the bone marrow for several weeks or months before it is eliminated. Recovered cats are immune to FeLV as evidenced by their resistance to subsequent challenge with virus.

Two factors influence the outcome of infection. The first is the age at which a cat is exposed to virus. Kittens up to 14 weeks of age are very susceptible and essentially all can be affected by virus challenge. Older kittens are more resistant so that by 16 weeks of age only about 20% can be infected. The other factor governing susceptibility is dose of virus: high doses lead to persistent viraemia while lower does immunise. The consequences of persistent infection are severe and essentially all viraemic cats die within four years of infection. By contrast recovered cats have a normal life span.

Rec very is due to an immune response t the virus. Only antibody responses have been examined in detail. However, it is clear that resistance to reinfection is mediated by virus neutralising antibodies which are directed at the envelope surface glycoprotein the virus, gp70. Attempts have been made to immunise cats against FeLV infection with vaccines containing gp70, some of which have been successful.

European patent specification EP 0377842 describes the production of replication-defective FeLV viral sequences. US patent 4,794,168 relates to the use of immunogens from FeLV envelope protein P15E in vaccines. European patent specifications 0247904, 0173997 and 0216564, International patent publication W085/02625 and US patent 4,789,702 all describe the use of the FeLV envelope protein gp70 or fragments thereof as antigens.

SUMMARY OF THE INVENTION

The present invention is based on the discovery that a fused recombinant protein comprising two regions of the FeLV molecule which are normally non-adjacent, when combined produced an immunogenic determinant capable of inducing anti FeLV antibodies on inoculation into cats. The immunised cats are resistant to challenge with live FeLV.

Thus, the present invention provides an immunogenic fused recombinant protein which comprises:

- a first pr tein which is feline l ukaemia virus (FeLV) gp70 VR1 protein or effective fragment thereof, having fused thereto;
- a second non-adjacent FeLV protein or effective fragment thereof different from the first protein and capable of stabilising the first protein, such that said fused recombinant protein is immunogenic and protective against FeLV infection.

It has been found that the use of gp70 VR1 protein alone, whilst producing antibodies, is not protective against FeLV infection.

It has however been found that the fusion of a second non-adjacent FeLV protein (or fragment thereof) may result in a fused recombinant protein which is both antigenic and protective against FeLV infection. The second protein is preferably also derived from gp70. In particular, the gp70 VR5 protein (or fragment thereof), which may not by itself produce antibodies, is capable of stabilising the first protein and producing a fused protein which surprisingly has immunogenic and protective properties.

Preferably, the second protein is fused to the carboxy terminus of VR1.

Usually, the fused recombinant protein will be free of glycosylation.

The fused recombinant protein, preferably further comprises an antigenic co-protein, such as Glutathione-S-transferase (GST) or beta-galactosidase.

These relatively large co-proteins solubilise the pr tein and facilitate production and purification thereof.

Moreover, the co-protein may act as an adjuvant in the sense of providing a generalised stimulation of the immune system. Preferably the co-protein is attached to the amino terminus of the fused protein.

Surprisingly, it has been found that co-proteins (such as GST or \$-galactosidase) may be attached to the amino terminus of the fused protein without adversely affecting immunogenicity thereof, and imparting desirable solubility and stability properties. However, it has been found that the attachment of other proteins to the carboxy terminus (whilst being useful in stabilising the fusion protein against degradation during preparation) may adversely affect its immunogenicity. It is postulated that this may be due to conformational changes brought about by the presence of the co-protein attached at the carboxy terminus. Therefore any co-protein attached at the carboxy terminus should be relatively small such as not to disturb the antigenic conformation of the fusion protein.

The present invention also provides a vaccine formulation which comprises the immunogenic fused recombinant protein together with a suitable carrier. Since the protein may be broken down in the stomach, it is preferably administered parenterally (including subcutaneous, intramuscular, intravenous, intradermal etc.

injecti n). Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampoules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use.

The vaccine formulation may also include adjuvant systems for enhancing the immunogenicity of the formulation, such as oil-in water systems and other systems known in the art.

The dosage will depend on the specific activity of the vaccine and can be readily determined by routine experimentation.

The present invention also provides a method of constructing a fused recombinant protein which comprises:

- introducing a DNA sequence coding for the first protein into a plasmid;
- opening the plasmid and introducing a DNA sequence coding for the second protein adjacent the first DNA sequence, and
 - expressing the recombinant protein.

In particular, the DNA sequences may be amplified using polymerase chain reaction (PCR). The amplified sequences may then be inserted one after another into adjacent positions in the plasmid. The plasmid may also include a gene coding for GST or similar co-protein.

The plasmid thus constructed may be expressed in any suitable bacteria, such as E. coli.

Whilst the invention has been described with reference to certain named proteins, such as gp70 VR1 and VR5 proteins, it is to be understood that this covers fragments of the naturally occurring protein and similar proteins (for example, having sequence homologies of 50% or greater) with additions, deletions or substitutions which do not substantially affect the immunogenic properties of the fused recombinant protein.

The second protein is one which does not normally lie adjacent to the first protein in the native FeLV gp70 protein. Non-adjacent means that it is not adjacent in the linearised amino acid sequence.

DESCRIPTION OF PREFERRED EMBODIMENTS

Embodiments of the present invention will now be described by way of example only with reference to the following figures and examples.

Figure 1 shows the plasmid pGEX-2T employed in the construction of the fused recombinant protein:

Figure 2 sh ws a terminal sequence including restriction sites;

Figure 3 shows the native FeLV gp70 VR1 sequence;

Figure 4 shows the native FeLV gp70 VR5 sequence; and

Figure 5 is a diagrammatic representation of the variable regions in the FeLV gp70 gene.

The following examples illustrate the construction and immunogenic properties of the fused recombinant protein GST-VR1-VR5.

Example 1

Construction of a Plasmid Expressing Recombinant gp70 Fusion Protein

The cloning vector used for these experiments is pGEX-2T (Smith & Johnston, Gene 67 (1988) 31-40). The FeLV gp70 gene fragment is cloned into this plasmid to produce a polypeptide fused to the -COOH terminus of the enzyme Glutathione-S-transferase (GST). Previous experiments have demonstrated that large fragments of gp70 (over 20kd) often result in problems of insolubility and degradation. In an attempt to circumvent these problems it was decided to express a range of smaller gp70 fragments. Furthermore, the regions chosen for expression are some of the "variable" regions of gp70, i.e. those regions showing the greatest variation between sub-groups A, B and C.

The regi n of the FeLV gp70 gene containing VR1 was

amplified using the following primers. The cloned envelope gene was the template used.

- 1) 5'GAG GGA TCC CTA CAT GTT GAC TTA TG 3' BamHI
- 2) 5'TAG GAA TTC ACA ATG TGT TCC CTT TG 3'

The PCR product was digested with the restriction enzymes BamHI and EcoRI the sites for which had been incorporated into the primers as shown above. There are no sites for these two restriction enzymes within the amplified fragment.

The digested PCR product was separated, by electrophoresis on an agarose gel, from any contaminating by products of the PCR reaction. The section of the agarose gel containing the DNA fragment was excised and the DNA removed by electroelution. After ethanol precipitation the DNA fragment was cloned into BamHI/EcoRI cut pGEX-2T.

After transformation of a recipient E.coli strain, clones harboring a recombinant pGEX-2T/VR1 plasmid were isolated.

The following primers were used to amplify a DNA fragment spanning the VR5 region:

- 5'GAC GAA TTC CAG GCT TTG TGC AAT AAG ACA CAA 3' ECORI
- 5' TAG GAA TTC GCA TGC GGT GAG TCC AGT GTT ACA 3' EcoRI SphI

Cloned FeLV gp70 gene was again the template used, the PCR product was digested with EcoRI prior to

purificati n on agarose. This PCR product was cloned into the pGEX-2T/VR1 plasmid described above. Clones carrying VR5 in the correct orientation with respect to VR1 were identified both by DNA sequence analysis and restriction enzyme analysis.

Example 2

Preparation of GST-VR1VR5 Recombinant Protein

The glutathione-S-transferase (GST) of Schistosoma japonicum is a small, very soluble enzyme which is tolerant of insertions at its carboxyl end. The gene (Sj26) encoding for this enzyme has been placed under the control of an inducible promoter and introduced into a plasmid together with an ampicillin resistance marker on it. A multiple restriction site has been introduced at the 3' end of Sj26, in tandem with a region coding for a thrombin cleavage site. The resulting plasmid (pGEX-2T) can produce large quantities of a fusion protein, which is extractable using affinity chromatography. The fusion protein is cleavable and the resulting peptide can be purified by repeating the affinity chromatography.

The following materials and methods are part of routine laboratory practice and can be found in "Molecular Cloning, A Laboratory Manual" second edition, Eds.

Sambrook, Fritsch and Maniatis. and also in "Current Protocols in Molecular Biology." Below is a detailed description of the materials and methods used.

Mat rials Enriched L Br th 100ml L Broth 1ml 2M glucose 1ml 1M mgCl₂

2000x IPTG
1g isopropyl D thio- galactopyranoside (IPTG)
8ml water make up to 10ml
Filter sterilise, dispense into aliquots of 0.5ml, freeze
at -20°C

1000x PMSF in isopropanol lmg/ml phenyl methyl sulphonyl fluoride

20% TX100 2ml TX100 8ml water Filter sterilise. Easier to handle at this concentration.

Glutathione/Tris 250ml 1.0M Tris 4.75ml water 30.73mg glutathione Filter sterilise.

a) Preparation of glutathione beads

A 15ml polypropylene tube is taken and 70 mg of dehydrated glutathione agarose beads are swollen with 10 mls of PBS/1% TX100 using gentle agitation for 30 minutes.

They are then precipitated at 2000 rpm for 5 minutes. The beads are resuspended in 5 mls of PBS to wash them, this procedure is repeated three times. (PBS/1%TX100 or PBS/tween are valid alternatives).

- b) Production of Fusion Protein
- Bacterial cells, transfected with a recombinant pGEX plasmid containing the DNA coding for the protein of interest, are cultured in 100 mls of enriched L Broth overnight in the presence of ampicillin.

- 2. The vernight culture is inoculated into 900 mls of enriched L Broth (with ampicillin). The bacterial suspension is grown up with vigorous shaking to an 0.D of approximately 0.6 (550nm) (about 2 hours) then IPTG is added to a final concentration of 50ug/ml. The culture is left for a further 90 minutes. Whilst inducing the culture, start to prepare the swollen, washed glutathione beads for use in stage 5 below.
- 3. The culture is then centrifuged at 6000 rpm for 15 minutes. The supernatant is discarded. The pellets are resuspended in 10 mls of PBS/1% TX100. 10ul of PMSF in isopropanol is added.
- 4. The concentrated bacterial suspension is sonicated in ice for 30 seconds and then centrifuged at 18000 rpm for 15 minutes. The supernatant is retained.
- 5. The beads are then resuspended in the supernatant from step 4 above and 20% TX100 added to a final concentration of 1% TX100. The mixture is gently agitated for 1 hour.
- 6. The mixture is centrifuged at 2000 rpm for 15 minutes and the beads washed 3 times as before. One ml of the solution of 20mM glutathione/50mM Tris is added to the beads and left for 30 minutes. The beads are again centrifuged out (the supernatant being retained) and washed once using another 1ml of the above glutathione solution. The washing is added to the supernatant.

7. The supernatant contains the fusion protein. An aliquot should be retained to verify this. The supernatant is then dialysed overnight in PBS to remove the excess glutathione. It may then be frozen at -200C.

The preparation exhibited a degree of degradation and/or incomplete synthesis. Nevertheless sufficient material from a 2 litre culture was obtained to conduct these experiments (4 mg. total protein).

Example 3

Vaccination

Since young kittens are most at risk of FeLV infection, the immunisation regime is designed to test the capacity of a vaccine to protect kittens at an early age, and to be compatible with the timing of vaccination against other feline viruses.

The route of challenge is by the oronasal route to simulate natural horizontal transmission of FeLV. A dose of virus is used which will cause persistent viraemia in over 90% of 14 week old kittens.

The schedule of vaccination and monitoring is in Table 1. The response of the kittens to vaccination challenge is given in Table 2.

The VR1-VR5 preparation elicited an immune response. A large proportion of the antibodies raised are directed against the GST portion of the recombinant

molecule as measured by ELISA. H wever, a competition ELISA using native GST and the recombinant protein also indicates that the immunised cats also responded to the FeLV component of the recombinant protein.

Western blot analysis using FeLV as antigen shows that three out of the four VR1-VR5 treated cats (59,61,62) produced easily detectable anti-gp70 antibodies and cat 60 gave a weak response on Western analysis. All controls were negative.

Virus neutralising antibodies were found in one cat prior to challenge and in a further 2 of the 4, four cats 6 weeks post challenge.

Three of the four vaccinated cats (59,61,62) were also found to be negative for virus isolation 3, 6 and 9 weeks post challenge, whilst all 4 control animals developed viraemia.

The cats responses to immunisation and challenge show that the VR1-VR5 preparation is able to produce an anti-gp70 antibody response and protect the animals from virus challenge.

TABLE 1

SCHEDULE OF VACCINATION AND SAMPLING

Age of Kitten	Week from	
(weeks)	challenge	Event
9	- 5	Blood sample
	·	Vaccination-1
12	-2	Blood sample
		Vaccination-2
14	0	Blood sample
		Challenge
17	3	Blood sample
20	6	Blood sample
23	9	Blood sample

TABLE 2 (VACCINATION RESULTS)

GROUP	CAT					TIME (WEEK	(S) REL	ATIVE TO	TIME (WEEKS) RELATIVE TO CHALLENGE				
	ſĿ,	*A* V	-13	N N	> «	12 N N	0 > K	3	A V B W	> «	3 2 2 3	> 4	2 6
VR-1+5	0 0 0 0 0 0 0 0	1 1 1 1	0000		1111	0000	1 1 1 1	0 0 0 4 + 1 + + + 3 + +	1+11	1 + 1 1	++++	1+11	0 0 128 128
CONT	68 69 07	1111	0000	-	1111	0000	1 1 1 1	1111	++++	++++	+1+1	++++	0000
A G P27 ar N G Virus E B anti-g		itigen in pla isolated fro neutralising p70 antibody	plag from ing ody	sma m plasma antibody (ELISA) (Western blot)	b10	(1)		·		•	·	•	•

SUBSTITUTE SHEET.

SEQUENCE LISTING (FIGURE 2)

SEQ ID NO

SEQUENCE TYPE : Nucleotide with derived protein sequence

SEQUENCE LENGTH : 51 base pairs

STRANDEDNESS

: single

: 1

TOPOLOGY

: linear

MOLECULAR TYPE : genomic DNA

ORIGINAL SOURCE

ORGANISM

: Feline leukaemia virus (FeLv)

USE

DEPOSIT

FeLV protein gp70 terminal sequence with

restriction sites

SEQUENCE LISTING (FIGURE 3)

SEQ ID NO : 2

SEQUENCE TYPE : 208 base pairs

STRANDEDNESS

: single

TOPOLOGY

: linear

MOLECULAR TYPE : genomic DNA

ORIGINAL SOURCE

ORGANISM

: Feline leukaemia virus (FeLV)

USE

: as vaccine in fused protein

DEPOSIT

FEATURES

FeLV protein gp70 VR1

SEQUENCE LISTING (FIGURE 4)

SEQ ID NO : 3

SEQUENCE TYPE : Nucleotide with derived protein sequence

SEQUENCE LENGTH : 42 base pairs

STRANDEDNESS : single

TOPOLOGY

: linear

MOLECULAR TYPE : genomic DNA

ORIGINAL SOURCE

ORGANISM

: Feline leukaemia virus (FeLV)

USE

: as vaccine in fused protein

DEPOSIT

FEATURES

FeLV protein gp70 VR5

CLAIMS

- 1. An immunogenic fused recombinant protein which comprises:
 - a first protein which is feline leukaemia virus (FeLV) gp70 VR1 protein or effective fragment thereof, having fused thereto;
 - a second non-adjacent FeLV protein or effective fragment thereof different from the first protein and capable of stabilising the first protein, such that said fused recombinant protein is immunogenic and protective against FeLV infection.
- 2. A fused protein according to claim 1 wherein said second protein is fused to the carboxy terminus of said first VR1 protein.
- 3. A fused protein according to any preceding claim wherein the second protein or fragment thereof is derived from FeLV gp70.
- 4. A fused protein according to claim 3 wherein the second protein is FeLV gp70 VR5 protein or effective fragment thereof.

- 5. A fused protein acc rding to any preceding claim which further comprises an antigenic co-protein attached thereto.
- 6. A fused protein according to claim 5 wherein the co-protein is attached to the amino terminus thereof.
- 7. A fused protein according to claim 5 or 6 wherein the co-protein is glutathione -S- transferase or 8-galactosidase.
- 8. A vaccine formulation which comprises the fusion protein of any preceding claim together with a pharmaceutically acceptable carrier.




Fig 2

CCA AAA TCG GAT CTG GTT CCG CGT GGA TCC CCG GGA ATT CAT CGT GAC TGA
Pro Lys Ser Asp Leu Val Pro Arg Gly Ser Pro Gly Ile His Arg Asp ***

Fig 3


3/5

FeLV VR1

Fig 4

FeLV VR5

CAG GCT TTG TGC AAT AAG ACA CAA CAG GGA CAT ACA GGG GCG Gln Ala Leu Cys Asn Lys Thr Gln Gln Gly His Thr Gly Ala CAC TAT CTA GCC GCC CCC AAC GGC ACC TAT TGG GCC TGT AAC His Tyr Leu Ala Ala Pro Asn Gly Thr Tyr Trp Ala Cys Asn ACT GGA CTC ACC Thr Gly Leu Thr



SUBSTITUTE SHEET

International Application No

	5 C12N15/49		National Classification and IPC); C12N15/62;	A61720/21
		0121137 1	5, C12N13/02;	A61K39/21
II. FIELDS	SEARCHED			
Classification	San Surrey	Minima	n Documentation Searched	
CHSINE	ou System		Classification Symbols	
Int.C1.	5 0	CO7K ; C12M	ı	
			hed other than Minimum Documentation connects are included in the Fields Searched	
III. DOCUM	ENTS CONSIDERED	TO BE RELEVANT		
Category °			appropriate, of the rejevant passages 12	Relevant to Claim N
		<u> </u>		
^	CORPORATI 2 Decembe cited in		BIOSCIENCE	1
A	WO,A,8 50 RESEARCH 7 Novembe see the w	4 871 (SCRIPPS C FOUNDATION) r 1985 hole document	LINIC AND	1
A	WO,A,8 50	the application _ 2 625 (CETUS COR	 PORATION)	1
	20 June 19 cited in s see the w	985 the application hole document —		
			-/-	-
"A" docum	neural to se ot Serrocites.	state of the art which is not	"I" later document published after or priority date and not in con- cited to understand the princi- invention	office with the application but ple or theory underlying the
"I." documentation of the communication of the comm	ent which may throw do its cited to establish the a or other special reason neat referring to an oral means	ubts on priority claim(s) or publication date of another a (as specified) disclosure, use, exhibition or as international filling date but	"X" document of particular releval cannot be considered anvel or involve an investive step "Y" document of particular releval cannot be considered to involve document is combined with or ments, such combination bein in the art.	cannot be considered to note; the claimed invention no an inventive step when the stor more other such docu-
later (min the priority date da	inel	"A" document member of the same	s patent family
	mai Completion of the L	ntermetional Security	Day of he w	
	27 AUGUST		Date of Mailing of this Internal 2 9 -09	mining Search Report
nternational 5	earching Authority		Signature of Authorized Office	•

International Application N

years, .		
	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No.
. 1	EP,A,O 173 997 (CHIRON CORPORATION)	1
	12 March 1986	•
	cited in the application	
	see the whole document	
-	ED A 0 202 240 (AMDAD CORROBATION LIMITED)	
	EP,A,O 293 249 (AMRAD CORPORATION LIMITED) 30 November 1988	7
- 1	see the whole document	
,P	PROCEEDINGS OF THE NATIONAL ACADEMY OF	1
	SCIENCES OF USA.	
1	vol. 89, no. 18, September 1992,	
ļ	WASHINGTON US pages 8457 - 8461	
	BROJATSCH, J. ET AL. 'Feline Leukemia	
	Virus subgroup C phenotype evolves through	
	distinct alterations near the N terminus	
- 1	of the envelope surface glycoprotein'	
ŀ	see the whole document	1
[DEUCEEDINGS OF THE NATIONAL ACADEMY OF	1.
	PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA.	1
İ	vol. 81, no. 12, June 1984, WASHINGTON US	
	pages: 3675 - 3679	,
1	NUNBERG, J.H. ET AL. 'Method to map	
	antigenic determinants recognized by	
	monocional antibodies: localization of a	
l	determinant of virus neutralization on the feline leukemia virus envelope protein	
1	I SING I CURCINIA Y I TUS PRIVATORA DYNTAIN	
	qp701	
	gp70'	
	gp70' see the whole document	
	gp70'	

ANNEX TO THE INTERNATIONAL SEARCH REPORT ON INTERNATIONAL PATENT APPLICATION NO.

GB 9300996 SA 74215

This amex lists the patent family members relating to the patent documents cited in the above-mentioned international search report.

The members are as contained in the European Patent Office EDP file on

The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

27/08/93

Patent document cited in search report	Publication date	Patr	ext family raber(s)	Publication date
EP-A-0247904	02-12-87	AU-B-	607427	07-03-91
		AU-A-	7547987	22-12-87
		JP-T-	1503273	09-11-89
		WO-A-	8707301	03-12-87
WO-A-8504871	07-11-85	US-A-	4663436	05-05-87
		AU-B-	589563	19-10-89
•		AU-A-	4354985	15-11-85
		CA-A-	1265648	06-02-90
		EP-A-	0179139	30-04-86
		JP-T-	61502126	25-09-86
*		US-A-	4794168	27-12-88
WO-A-8502625	20-06-85	US-A-	4701416	20-10-87
		AU-B-	577557	29-09-88
		AU-A-	3678184	26-06-85
		DE-A-	3484991	02-10-91
		EP-A.B	0164403	18-12-85
		JP-A-	2238884	21-09-90
		JP-T-	61500662	10-04-86
		-A-2U	4789702	06-12-88
EP-A-0173997	12-03-86	US-A-	4876089	24-10-89
		US-A-	5152982	06-10-92
EP-A-0293249	30-11-88	AU-B-	607511	07-03-91
		AU-A-	1793288	21-12-88
		MO-Y-	8809372	01-12-88
		DE-A-	3873989	01-10-92
		JP-T-	1503441	22-11-89

PORM FOOD

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82